

(١)

1-

(b) 3 or 2



2-


(b) The Curve of the function  $f$  has  
a local minimum value at  $x=3$



3-

$$\therefore y = ax^b$$

$$\therefore \frac{dy}{dt} = abx^{b-1} \frac{dx}{dt}$$



$$\frac{dy}{dt} = \frac{abx^b}{x} \cdot \frac{dx}{dt} \quad \text{Multiply by } \frac{1}{y}$$


$$\therefore \frac{1}{y} \cdot \frac{dy}{dt} = \frac{1}{ax^b} \cdot \cancel{abx^b} \cdot \frac{dx}{dt} \quad \frac{1}{2}$$

$$\frac{1}{y} \cdot \frac{dy}{dx} = \frac{b}{x} \cdot \frac{dx}{dt} \quad \frac{1}{2}$$

4-

$$\begin{aligned}
 V &= \pi \int_{-2}^2 y^2 dx \\
 &= 2\pi \int_0^2 (x^2 + 2)^2 dx \\
 &= 2\pi \int_0^2 (x^4 + 4x^2 + 4) dx \\
 &= 2\pi \left[ \frac{1}{5}x^5 + \frac{4}{3}x^3 + 4x \right]_0^2 \\
 &= 2\pi \left[ \frac{32}{5} + \frac{32}{3} + 8 \right] \\
 &= \frac{752}{15}\pi
 \end{aligned}$$

Volume unit.



(تراعى الحلول الأخرى)

٣

5-

(b)  $\frac{1}{3} \ln 2$

6-

(d) 2

7-

$\therefore$  The area of the circular sector = 4

$$\therefore \frac{1}{2} r L = 4 \Rightarrow L = \frac{8}{r}$$


let the perimeter of the sector = y

$$\therefore y = 2r + L \Rightarrow y = 2r + \frac{8}{r}$$

$$\therefore \frac{dy}{dr} = 2 - \frac{8}{r^2}$$

$$\text{at } \frac{dy}{dr} = 0 \Rightarrow \frac{8}{r^2} = 2 \Rightarrow r^2 = 4 \Rightarrow r = 2 \text{ cm}$$

Discuss the sign of the function:



at  $r = 2 \text{ cm}$

The perimeter is minimum as possible

$$\therefore L = \frac{8}{2} = 4, \theta^{\text{rad}} = \frac{L}{r} = \frac{4}{2} = 2^{\text{rad.}}$$

٤

8-

To determine the intersection point:


$$\text{let } y_2 = y_1 \Rightarrow x+2 = 4-x^2 \\ \therefore x^2 + x - 2 = 0$$

$$(x+2)(x-1)=0$$

$$x = -2$$

or

$$x = 1$$




$$\text{Area} = \int_{-2}^1 [(4-x^2) - (x+2)] dx$$



$$= \int_{-2}^1 [4 - x^2 - x - 2] dx$$

$$= \int_{-2}^1 [-x^2 - x + 2] dx$$

$$= \left[ -\frac{1}{3}x^3 - \frac{1}{2}x^2 + 2x \right]_{-2}^1$$



$$= \left( -\frac{1}{3} - \frac{1}{2} + 2 \right) - \left( \frac{8}{3} - 2 - 4 \right)$$

$$= \frac{9}{2} \text{ area unit}$$



(تراعى الحلول الأخرى)

٥

9-

(b)  $\log_a b$



10-

(c) ٤



٦

11-

$$\text{Q} \int x^3 (x^2 + 1)^6 dx$$

$$\text{let } y = x^2 + 1 \Rightarrow x^2 = y - 1$$

$$\therefore 2x dx = dy \Rightarrow dx = \frac{dy}{2x} \quad \boxed{\frac{1}{2}}$$

$$\therefore \int x^2 \cdot x (x^2 + 1)^6 dx \quad \boxed{\frac{1}{2}}$$

$$= \int (y-1) \cdot x \cdot y^6 \cdot \frac{dy}{2x} \quad \boxed{\frac{1}{2}}$$

$$= \int \left( \frac{1}{2} y^7 - \frac{1}{2} y^6 \right) dy \quad \boxed{\frac{1}{2}}$$

$$= \frac{1}{16} y^8 - \frac{1}{14} y^7 + C \quad \boxed{\frac{1}{2}}$$

$$= \frac{1}{16} (x^2 + 1)^8 - \frac{1}{14} (x^2 + 1)^7 + C \quad \boxed{\frac{1}{2}}$$

$$\text{B} \int (x-3) e^{2x} dx$$

$$= (x-3) \left( \frac{1}{2} e^{2x} \right) - \int \frac{1}{2} e^{2x} dx \quad \begin{array}{l} \text{let } u = x-3 \\ dv = e^{2x} dx \\ du = dx \\ v = \frac{1}{2} e^{2x} \end{array}$$

$$= \frac{(x-3)}{2} e^{2x} - \left[ \frac{1}{2} \times \frac{1}{2} e^{2x} \right] + C \quad \boxed{\frac{1}{2}}$$

$$= \frac{(x-3)}{2} e^{2x} - \frac{1}{4} e^{2x} + C \quad \boxed{\frac{1}{2}}$$


(تراعى الحلول الأخرى)

٧

12-

(a)  $-\ln |\cos \varphi| + C$  

13-

(b) Zero 

٨

14-

a)  $f(x) = x^3 - 3x - 2$

$$f'(x) = 3x^2 - 3, f''(x) = 6x$$

$$\text{at } f'(x) = 0 \Rightarrow 3x^2 - 3 = 0$$

$$3(x-1)(x+1) = 0$$

$$\therefore x = 1 \quad \text{or} \quad x = -1$$



From the Discuss of the sign of the function:

$f(x)$  has a local max.



value at  $x = -1$

which is  $f(-1) = \text{zero}$



$f(x)$  has a local min. value at  $x = 1$

which is  $f(1) = -4$




at  $f''(x) = 0 \Rightarrow 6x = 0 \Rightarrow x = 0$





From the discuss of the sign of the function:

$f(x)$  has an inflection



point at  $x = 0$

which is  $(0, f(0))$



b)  $f(x) = x(x^2 - 12) = x^3 - 12x$

$$\therefore f'(x) = 3x^2 - 12$$

$$\text{at } f'(x) = 0 \Rightarrow 3(x^2 - 4) = 0 \Rightarrow x = \pm 2$$

$$2 \in [-1, 4] \quad \text{and} \quad -2 \notin [-1, 4]$$


$$f(-1) = 11, f(2) = -16, f(4) = 16$$



absolute min. value



absolute max. value



(تراعى الحلول الأخرى)

٩

15-

(a)  $-50$



16-

(b)  $]-\infty, 0]$




17-

$$x = \sec \theta, \quad y = \tan \theta$$

$$\text{at } \theta = \frac{\pi}{6} \Rightarrow x = \sec \frac{\pi}{6} \Rightarrow x = \frac{2\sqrt{3}}{3}$$

$$\Rightarrow y = \tan \frac{\pi}{6} \Rightarrow y = \frac{\sqrt{3}}{3}$$

$$\left( \frac{2\sqrt{3}}{3}, \frac{\sqrt{3}}{3} \right) \in \text{Curve}$$



$$\therefore \frac{dx}{d\theta} = \sec \theta \cdot \tan \theta, \quad \frac{dy}{d\theta} = \sec^2 \theta$$



$$\therefore \frac{dy}{dx} = \frac{\sec^2 \theta}{\sec \theta \cdot \tan \theta} = \frac{\sec \theta}{\tan \theta} = \csc \theta$$




$$\text{at } \theta = \frac{\pi}{6} \quad \text{The slope} = 2$$



The equation of the tangent:

$$y - \frac{\sqrt{3}}{3} = 2 \left( x - \frac{2\sqrt{3}}{3} \right)$$



The equation of the normal:

$$y - \frac{\sqrt{3}}{3} = -\frac{1}{2} \left( x - \frac{2\sqrt{3}}{3} \right)$$



(١٠)

١٨-

$$\sin y + \cos 2x = 0$$

$$\cos y \frac{dy}{dx} - 2 \sin 2x = 0$$

$$\cos y \frac{d^2y}{dx^2} + (-\sin y \frac{dy}{dx})(\frac{dy}{dx}) - 4 \cos 2x = 0$$



$$\cos y \frac{d^2y}{dx^2} - \sin y \left(\frac{dy}{dx}\right)^2 = 4 \cos 2x \quad \div (\cos y)$$

$$\frac{d^2y}{dx^2} - \left(\frac{dy}{dx}\right)^2 \tan y = 4 \cos 2x \sec y .$$



(تراعى الحلول الأخرى)

(انتهت الإجابة وتراعى الحلول الأخرى)